

Datasheet: STAR117D405GA

Description:	GOAT ANTI MOUSE IgG (H/L):DyLight®405 (MULTI SPECIES ADSORBED)
Specificity:	IgG (H/L)
Format:	DyLight®405
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/25 - 1/250
Immunofluorescence				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

ed to DyLight®405 - lid	quid	
Purified IgG conjugated to DyLight®405 - liquid		
Excitation Max (nm)	Emission Max (nm)	
400	420	
	` '	

Antiserum Preparation Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen. Purified IgG prepared by affinity chromatography

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Whole mouse IgG
Fortament Databases	

External Database Links

UniProt:

P01837 Related reagents
P01869 Related reagents

P01867	Related reagents
P01864	Related reagents
P01843	Related reagents
P01865	Related reagents
P01844	Related reagents
P01868	Related reagents
P01724	Related reagents
P03987	Related reagents
P01863	Related reagents
P01845	Related reagents

Entrez Gene:

<u>16071</u>	lgk-C	Related reagents
<u>16017</u>	lghg1	Related reagents
<u>16016</u>	lghg2b	Related reagents
380793	lgh-1a	Related reagents
380793	lgh-1a	Related reagents
<u>433053</u>	LOC433053	Related reagents
<u>16017</u>	lghg1	Related reagents
<u>16142</u>	lglv1	Related reagents
<u>110786</u>	lglc2	Related reagents
110787	lalo2	Doloted research
110707	lglc3	Related reagents
380793	lgh-1a	Related reagents

Synonyms

lgh-4

Specificity

Goat anti Mouse IgG antibody recognizes mouse IgG and light chains common to other mouse immunoglobulin classes.

Goat anti Mouse IgG has been cross-adsorbed using human, bovine, porcine, equine, lapine and chicken immunoabsorbants to remove cross-reactive antibodies. Less than 0.1% cross reactivity was detected to human, bovine, porcine, equine, caprine, lapine and chicken IgG by immunoelectrophoresis and ELISA.

Goat anti Mouse IgG antibody is highly recommended for use as a secondary antibody with human and veterinary samples. Goat anti Mouse IgG antibody has been used successfully as a secondary detection reagent in combination with mouse clone $\underline{\text{CC327}}$ for the detection of $\overline{\text{TNF}\alpha}$ and mouse clone $\underline{8M6}$ for the detection of interleukin-8 in bovine respiritory syncitial virus infected, neonatal ovine lung tissue by immunohistochemistry ($\underline{\text{Redondo } et al. 2013}$).

Flow Cytometry

Use 50 ul of the suggested working dilution to label 1x10⁶ cells in 100ul

References

- 1. Abdala-Valencia, H. *et al.* (2012) Vitamin E isoforms differentially regulate intercellular adhesion molecule-1 activation of PKCα in human microvascular endothelial cells. PLoS One. 7: e41054.
- 2. Redondo, E. *et al.* (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. <u>J Comp Pathol. 150 (4): 434-48.</u>
- 3. Banerjee, K. *et al.* (2012) Occluding the mannose moieties on human immunodeficiency virus type 1 gp120 with griffithsin improves the antibody responses to both proteins in mice. <u>AIDS Res</u>

Hum Retroviruses. 28 (2): 206-14.

- 4. Singh, S.M. et al. (2016) Characterization of Immune Responses to an Inactivated Avian Influenza Virus Vaccine Adjuvanted with Nanoparticles Containing CpG ODN. Viral Immunol. Apr 14. [Epub ahead of print]
- 5. Iwaszko-Simonik, A. et al. (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). Vet Immunol Immunopathol. 164 (1-2): 87-92.
- 6. Askari, N. et al. (2015) Tetracycline-regulated expression of OLIG2 gene in human dental pulp stem cells lead to mouse sciatic nerve regeneration upon transplantation. Neuroscience. 305: 197-208.
- 7. Topoluk, N. et al. (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. Am J Sports Med.: 363546517706138.
- 8. Alimolaei, M. et al. (2017) A Recombinant Probiotic, Lactobacillus casei, Expressing the Clostridium perfringens α-toxoid, as an Orally Vaccine Candidate Against Gas Gangrene and Necrotic Enteritis. Probiotics Antimicrob Proteins. Apr 11 [Epub ahead of print].
- 9. Schmidli, M.R. et al. (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. BMC Vet Res. 14 (1): 161.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
Acknowledgements	DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

North & South Tel: +1 800 265 7376 America

Regulatory

Fax: +1 919 878 3751

Worldwide

For research purposes only

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M314241:180412'

Printed on 31 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint